

On the impossibility of measuring the general relativistic part of the terrestrial acceleration of gravity with superconducting gravimeters

Lorenzo Iorio
*Viale Unità di Italia 68, 70125
Bari, Italy
e-mail: lorenzo.iorio@libero.it*

Abstract

In this paper we very preliminarily investigate the possibility of measuring the post-Newtonian general relativistic gravitoelectric and gravitomagnetic components of the acceleration of gravity on the Earth, in continuous regime, with two absolute measurements at the equator and the south pole with superconducting gravimeters. The magnitudes of such relativistic effects are $10^{-10} \text{ m s}^{-2}$ and $10^{-11} \text{ m s}^{-2}$, respectively. Unfortunately, the present-day uncertainties in the Earth's geodetic parameters which enter the classical Newtonian terms induce systematic errors 1-2 orders of magnitude larger than the relativistic ones. Moreover, a $\sim 1 \text{ ngal}$ sensitivity can be reached by the currently available superconducting gravimeters, but only for relative measurements.

1 Introduction

The classical test-bed of the Einstein's General Theory of Relativity (GTR), in its linearized weak-field and slow-motion approximation [1] valid in the Earth's neighborhood and throughout the Solar System, has always been represented by the motion of test particles and electromagnetic waves in the gravitational field of massive astronomical bodies [2]. This is so due to the extreme smallness of the relativistic effects with respect to the much larger Newtonian features of gravity.

However, in the more or less recent past various Earth-based laboratory experiments [3, 4, 5, 6, 7, 8, 9] have been proposed, mainly to detect the elusive gravitomagnetic part of the gravitational field of the Earth induced by its proper angular momentum (see Section 2). Up to now, they have not been performed, mainly due to the extreme difficulty both to reach the

necessary high sensitivity and to cope with the numerous and much larger classical noising effects of natural and cultural origin.

In regard to the possibility of testing GTR with terrestrial geodetic techniques [1], gravimetry can allow to penetrate into the relativistic regime. Among the most sensitive apparatus there are the superconducting gravimeters (SG) [10]. Their most remarkable quality is their high immunity to the environmental temperature variations, that generally determine variations of the elastic constant of the mechanical spring which, indeed, is replaced by a magnetic levitation of a superconducting sphere in the magnetic field of superconducting, persistent current coils. The goal is to utilize the high stability of such currents to create a perfectly stable spring. The magnetic levitation is designed to provide independent adjustment of the total levitating force and the force gradient so that it can support the full weight of the sphere and still yield a large displacement for a small change in gravity. Recent developments in such a field have pushed the accuracy of SG down to $10^{-12}g = 10^{-11} \text{ m s}^{-2} = 1 \text{ ngal}$ [11] for relative measurements. For some SG performed tests of certain non-relativistic preferred-frame and preferred-locations effects, in the context of the Parameterized Post Newtonian (PPN) formalism, see Section 8.2 of [12]. However, it must also be noted that there are great difficulties to use SG in absolute measurements due to calibration issues.

2 The post-Newtonian general relativistic components of the acceleration of gravity

The acceleration experienced by a test mass in the gravitational field of a central body, to the post-Newtonian $\mathcal{O}(c^{-2})$ level of GTR in its linearized weak-field and slow-motion approximation, is represented by a gravitoelectric component, due to the Schwarzschild part [13] of the spacetime metric, and a gravitomagnetic component, due to the Lense-Thirring part [14] of the spacetime metric. We can write it as $\mathbf{a} \equiv \mathbf{a}_{\text{Newton}} + \mathbf{a}_{\text{Schwarzschild}} + \mathbf{a}_{\text{Lense-Thirring}}$ with [1]

$$\begin{cases} \mathbf{a}_{\text{Newton}} &= -\frac{GM}{r^3}\mathbf{r}, \\ \mathbf{a}_{\text{Schwarzschild}} &= \frac{GM}{c^2 r^3} \left\{ \left[\frac{4GM}{r} - v^2 \right] \mathbf{r} + 4(\mathbf{r} \cdot \mathbf{v})\mathbf{v} \right\}, \\ \mathbf{a}_{\text{Lense-Thirring}} &= \frac{2G}{c^2 r^5} \left[3(\mathbf{r} \cdot \mathbf{J})\mathbf{r} \times \mathbf{v} + r^2 \mathbf{v} \times \mathbf{J} \right]. \end{cases} \quad (1)$$

G is the Newtonian gravitational constant, c is the speed of light in vacuum, M is the mass of the central body, $\mathbf{J}$ is its proper angular momentum¹, $\mathbf{r}$ and $\mathbf{v}$ are the position and velocity vectors, respectively, of the test mass. It is assumed that the origin of the adopted inertial reference frame is located at the centre of mass of the body of mass M .

3 Absolute measurements at the pole and the equator

We will now explore the possibility of measuring the gravitoelectric and gravitomagnetic components of the terrestrial gravitational field by measuring the acceleration of gravity at the equator and at the South Pole and making a cross-check between these two absolute measurements. To this aim, let us, now, consider a test body fixed on the Earth's surface, so that $\mathbf{v} = \boldsymbol{\omega} \times R\hat{\mathbf{r}}$ and $v = \omega R \cos \lambda$ where $\boldsymbol{\omega} = \omega \mathbf{k}$ is the Earth's angular velocity vector, $\hat{\mathbf{r}}$ is a unit vector from the Earth's centre to the location of the test mass, λ is the geocentric latitude and R is the Earth's radius at the location of the test body. From (1) it is easy to see that at the equator ($\mathbf{r} \cdot \mathbf{J} = 0$) the Lense-Thirring acceleration is entirely radial and directed outward, as the centrifugal acceleration, while at the South Pole it is absent ($\mathbf{v} = 0$). The gravitoelectric Schwarzschild component is always directed radially because at the equator $\mathbf{r} \cdot \mathbf{v} = 0$ and at the poles $\mathbf{v} = 0$. Then, the acceleration of gravity can be written as

$$\begin{cases} g_{\text{eq}} &= \frac{GM}{R_{\text{eq}}^2} - \omega^2 R_{\text{eq}} + \frac{GM}{c^2 R_{\text{eq}}^3} (4GM - \omega^2 R_{\text{eq}}^3) - \frac{2GJ\omega}{c^2 R_{\text{eq}}^2}, \\ g_{\text{pol}} &= \frac{GM}{R_{\text{pol}}^2} + \frac{4(GM)^2}{c^2 R_{\text{pol}}^3}, \end{cases} \quad (2)$$

where the difference between the Earth's radius at the poles and at the equator is related to the Earth's flattening f by

$$\frac{R_{\text{eq}} - R_{\text{pol}}}{R_{\text{eq}}} \equiv f. \quad (3)$$

Then, we can write for $\Delta g \equiv g_{\text{pol}} - g_{\text{eq}}$

$$\Delta g \simeq \frac{2GM}{R_{\text{eq}}^2} \frac{f}{(1-f)^2} + \frac{12(GM)^2}{c^2 R_{\text{eq}}^3} \frac{f}{(1-f)^3} + \omega^2 \left(R_{\text{eq}} + \frac{GM}{c^2} \right) + \frac{2GJ\omega}{c^2 R_{\text{eq}}^2}. \quad (4)$$

Table 1: Relevant geodetic parameters.

Gravitational constant G ($\text{kg}^{-1} \text{ m}^3 \text{ s}^{-2}$) [15]	$6.673 \times 10^{-11} \pm 1 \times 10^{-13}$
Speed of light c (m s^{-1}) [15]	2.99792458×10^8
Earth's GM ($\text{m}^3 \text{ s}^{-2}$) [16]	$3.986004418 \times 10^{14} \pm 8 \times 10^5$
Earth's mean equatorial radius R_{eq} (m) [16]	$6.3781366 \times 10^6 \pm 1 \times 10^{-1}$
Earth's angular momentum J ($\text{kg m}^2 \text{ s}^{-1}$) [17]	$5.85386532242 \times 10^{33}$
Earth's flattening f [16]	$3.352 \times 10^{-3} \pm 1.1 \times 10^{-10}$
Earth's angular speed ω (rad s^{-1}) [16]	$7.292115 \times 10^{-5} \pm 1 \times 10^{-12}$

Now we will evaluate the magnitude of the various terms entering (4) in order to compare them with the present-day available sensitivity of SG. Then, we will also evaluate the systematic errors induced by the classical terms of (4) due to the uncertainties in the various geodetic parameters of the Earth. Of course, there are also various classical time-dependent competing surface gravity effects² spanning a wide range of periodicities from 1 s to more than 1 year and magnitudes up to $1 - 10 \mu\text{gal} = 10^{-5} - 10^{-4} \text{ m s}^{-2}$ which would act as systematic bias³. They should be accounted for in a detailed error budget, which is, however, beyond the scope of the present paper.

3.1 The Lense-Thirring component

By using the values of Table 1 it turns out that the Lense-Thirring component $\Delta g_{\text{LT}} \equiv 2GJ\omega/c^2 R_{\text{eq}}^2$ of (4) amounts to $1.5 \times 10^{-11} \text{ m s}^{-2}$, i.e. $\sim 1 \text{ ngal}$.

3.1.1 The systematic errors

In order to obtain Δg_{LT} we should subtract the first three terms of (4) from the measured Δg . This can only be done if the uncertainty of the terms

¹For a homogeneous spherical body of mass M , radius R spinning at a rate ω $J = (2/5)MR^2\omega$; for the Earth the factor $2/5 = 0.4$ is replaced by 0.330841 [17].

²They are, e.g., the ocean noise, seismic and normal modes, slow and silent earthquakes, secular deformations.

³It must be noted that the investigated measurement would be in continuous regime.

to be subtracted is smaller than the predicted value of the Lense-Thirring component.

The residual Schwarzschild component would not pose problems. Indeed, its nominal value is

$$\Delta g_{\text{Schwarzschild}} = \frac{12(GM)^2}{c^2 R_{\text{eq}}^3} \frac{f}{(1-f)^3} + \frac{GM}{c^2} \omega^2 = 3.00 \times 10^{-10} \text{ m s}^{-2}. \quad (5)$$

According to the uncertainties released in Table 1 it turns out that $\delta(\Delta g_{\text{Schwarzschild}}) \ll \Delta g_{\text{LT}}$. The problems come from the classical terms. Indeed,

$$\Delta g_{\text{Newton}} = \frac{2GM}{R_{\text{eq}}^2} \frac{f}{(1-f)^2} = 6.6130308646 \times 10^{-2} \text{ m s}^{-2}, \quad (6)$$

which is nine orders of magnitude larger than the gravitomagnetic term. The errors induced by the uncertainty in f and the Earth's mean equatorial radius R_{eq} are the largest ones and amount to

$$\delta(\Delta g_{\text{Newton}})|_f = \frac{2GM}{R_{\text{eq}}^2} \frac{(1+f)}{(1-f)^3} \delta f = 2.184 \times 10^{-9} \text{ m s}^{-2} \quad (7)$$

and

$$\delta(\Delta g_{\text{Newton}})|_{R_{\text{eq}}} = \frac{4GMf}{R_{\text{eq}}^3 (1-f)^2} \delta R_{\text{eq}} = 2.073 \times 10^{-9} \text{ m s}^{-2}, \quad (8)$$

respectively. The centrifugal component amounts to

$$\Delta g_{\text{centrifugal}} = \omega^2 R_{\text{eq}} = 3.3906727993 \times 10^{-2} \text{ m s}^{-2}, \quad (9)$$

with an error of $\delta(\Delta g_{\text{centrifugal}}) = 1.48 \times 10^{-9} \text{ m s}^{-2}$. The total error in the classical part of Δg is, thus

$$\delta(\Delta g)_{\text{total}}^{\text{class}} \leq 5.737 \times 10^{-9} \text{ m s}^{-2}; \quad (10)$$

it is two orders of magnitude larger than Δg_{LT} . This rules out the possibility of measuring the gravitomagnetic Lense-Thirring component of the acceleration of gravity with the present-day SG.

3.2 The Schwarzschild component

From (2) it can be noted that an absolute measurement at the South Pole could allow to detect the gravitoelectric Schwarzschild component of the acceleration of gravity. Indeed, it turns out that

$$g_{\text{pol}}^{\text{Schwarzschild}} = \frac{4(GM)^2}{c^2 R_{\text{eq}}^3 (1-f)^3} = 2.752 \times 10^{-8} \text{ m s}^{-2}. \quad (11)$$

3.2.1 The systematic errors

On the other hand, the errors in the classical part are $\delta g_{\text{pol}}^{\text{class}}|_{GM} = 1.979 \times 10^{-8} \text{ m s}^{-2}$, $\delta g_{\text{pol}}^{\text{class}}|_{R_{\text{eq}}} = 3.0724 \times 10^{-7} \text{ m s}^{-2}$, $\delta g_{\text{pol}}^{\text{class}}|_f = 1.97 \times 10^{-9} \text{ m s}^{-2}$. The total systematic error would then be

$$\delta g_{\text{pol}}^{\text{class}}|_{\text{total}} = 3.29 \times 10^{-7} \text{ m s}^{-2}, \quad (12)$$

i.e. one order of magnitude larger than $g_{\text{pol}}^{\text{Schwarzschild}}$. Note that (5) and (10) rule also out the possibility of a measurement of the gravitoelectric component of the acceleration of gravity between the pole and the equator by one order of magnitude.

4 Discussion and conclusions

In this paper we have preliminarily investigated the feasibility of an experiment aimed at the measurement of the general relativistic gravitoelectric and gravitomagnetic components of the terrestrial acceleration of gravity at the equator and at the pole. It turns out that they are quite small; the difference between the gravitoelectric accelerations in the two places amounts to $3 \times 10^{-10} \text{ m s}^{-2}$ and the difference of the gravitomagnetic accelerations is $1.5 \times 10^{-11} \text{ m s}^{-2}$, i.e. 10 ngal and 1 ngal, respectively. Such absolute measurements could be done, in principle, with SG. However, it must be noted that the greatest difficulty in the use of the SG as an absolute instrument is connected with its calibration. Another non-trivial difficulty is that the effects to be measured are in continuous. Moreover, the present-day sensitivity of SG is just of the order of ngal, but only for relative measurements. In regard to systematic bias, the current uncertainties in the Earth's geodetic parameters, which enter the classical Newtonian terms to be subtracted, induce errors 1-2 orders of magnitude larger than the relativistic ones.

Acknowledgements

I thank the anonymous referees for their helpful and important comments which greatly improved the present paper.

References

- [1] Soffel, M.H., 1989, *Relativity in Astrometry, Celestial Mechanics and Geodesy*, (Springer, Berlin).

- [2] Will, C.M., 2001, *Living Rev. Relativ.* **4**, 4
<http://www.livingreviews.org/Articles/Volume4/2001-4will>
- [3] V.B. Braginsky, V.B., Polnarev, A.G., and Thorne, K.S., 1984, *Phys. Rev. Lett.*, **53**, 863.
- [4] Pippard, A.B., 1988, *Proc. R. Soc. London A*, **420**, 81.
- [5] Braginsky, V.B., Caves, C., and Thorne, K.S., 1977, *Phys. Rev. D*, **15**, 2047.
- [6] Cerdonio, M., Prodi, G.A., and Vitale, S., 1988, *Gen. Rel. Grav.*, **20**, 83.
- [7] Tartaglia, A., and Ruggiero, M.L. 2002, *Gen. Rel. Grav.*, **34**, 1371.
- [8] Stedman, G.E., Schreiber, K.U., and Bilger, H.R., 2003, *Class. Quantum Grav.*, **20**, 2527.
- [9] Iorio, L., 2003, *Class. Quantum Grav.*, **20**, L5.
- [10] Goodkind, J.M., 1999, *Review of Scientific Instruments*, **70**, 4131.
- [11] Francis, O., and van Dam, T., 2002, *Metrologia*, **39**, 485.
- [12] Will, C.M., 1993, *Theory and experiment in gravitational physics*, revised edition, (Cambridge University Press, Cambridge).
- [13] Schwarzschild, K., 1916, *Sitzungsberichte der Königlich Preussischen Akademie der Wissenschaften zu Berlin, Phys.-Math. Klasse*, 189 translated and discussed by Antoci, S., 2003, *Gen. Rel. Grav.*, **35**, 951.
- [14] Lense, J., and Thirring, H., 1918, *Phys. Z.* **19**, 156 translated and discussed by Mashhoon, B., Hehl, F. W., and Theiss, D. S., 1984, *Gen. Rel. Grav.*, **16**, 711.
- [15] Mohr, P.J., and Taylor, B.N., 1999, *J. Phys. Chem. Ref. Data*, **28**, 6, p. 1713.
- [16] Groten, E., 1999, *Report of the IAG Special Commission SC3, Fundamental Constants*, XXII IAG General Assembly.
- [17] McCarthy, D. D., and Petit, G., 2004, *IERS conventions (2003)* IERS Technical Note 23 (Verlag des Bundesamtes für Kartographie und Geodäsie, Frankfurt am Main).